



MARKING SCHEME

**LEVEL 2 CERTIFICATE IN ADDITIONAL
MATHEMATICS
9550/01**

SUMMER 2017

INTRODUCTION

This marking scheme was used by WJEC for the 2017 examination. It was finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conference was held shortly after the paper was taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conference, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about this marking scheme.

LEVEL 2 CERTIFICATE IN ADDITIONAL MATHEMATICS

MARK SCHEME - SUMMER 2017

| | Additional Mathematics Summer 2017 | | Final Version |
|---|---|---|--|
| 1 | $(5x + 3)(4x - 1)$ $-3/5$ and $1/4$ | B2 B2 4 | $B1 (5x \dots 3)(4x \dots 1)$ or $(5x \dots 1)(4x \dots 3)$ or $5x(4x - 1) + 3(4x - 1)$ or $4x(5x + 3) - 1(5x + 3)$, or $(20x + 12)(x - 1/4)$ B0 for $(5x + 3)(20x - 5)$ Ignore sight of “=0” Must be from factorising, do not accept use of quadratic formula followed by ‘factorising’. MUST FT for their factors FT for ‘their factors’ equivalent difficulty not leading to whole number solutions. B1 for each answer |
| 2 | (a) $70x^9 - 5 (+0)$ (b) $-12x^{-13}$ or $-12/x^{13}$ (c) $3/8x^{-5/8}$ or equivalent (d) $-4x^{-5}$ or $-4/x^5$ | B3 B1 B1 B1 6 | $B1$ for $70x^9$ (not $10 \times 7x^9$), $B1$ for -5 and $B1$ for $+0$ (or blank) provided at least one other mark awarded. If B3 penalise further incorrect working -1 CAO , although ISW. Index needs to be simplified. CAO , although ISW. Index needs to be simplified. CAO , although ISW. Index needs to be simplified. Penalise including ‘+c’ -1 only |
| 3 | $(x + 11)^2 (\pm \dots)$ or $(x + 22/2)^2 (\pm \dots)$ (Minimum value at $x =$) -11 (Minimum value is) (+) 2 | M1 A1 A1 3 | Ignore ‘their ($\pm \dots$)’ or ‘=0’ Do not accept method $dy/dx = 2x + 22$ CAO . Must be from sight of completing the square CAO . Must be from sight of completing the square |
| 4 | (a) $2(-4)^3 - 5(-4)^2 + 8(-4) - 6 (= -128 - 80 - 32 - 6)$ -246 (b)(i) Substitute $x = 2$ Showing $f(2) = 0$ (ii) $(x - 2)(x^2 + bx + c)$ or intention to divide by $(x - 2)$ with x^2 shown $(x - 2) (x^2 + 11x + 30)$ $(x - 2) (x + 5)(x + 6)$ | M1 A1 M1 A1 M1 A2 A1 8 | Or division method giving $2x^2 - 13x \dots$ Or division method giving $x^2 + 11x \dots$ Accept sight of substitution with ‘=0’ shown Or equivalent ie. $X^2 + 11x - 60$ with no remainder If any values are inserted at least 1 needs to be correct, appropriate sight of $+11x$ or $+30$ implies M1 (and A1 to follow) $A1$ for $(+11x$ or $(+30)$ Or use of factor theorem $A1 (x + 5)$, $A1 (x + 6)$ CAO , but ignore sight of “=0”, ISW |

| | Additional Mathematics Summer 2017 | | Final Version |
|---|---|---|---|
| 6 | <p>(a) $(FG^2 =) (20 - 10)^2 + (8 - -4)^2 (=10^2 + 12^2)$</p> $FG = \sqrt{244}$ $= 2\sqrt{61}$ <p>(b) Gradient FG $(20 - 10) / (8 - -4)$ $= 10/12 (= 5/6)$ Gradient perpendicular $-12/10 (= -6/5)$</p> <p>$(8 + -4)/2, (20 + 10)/2$ Mid point FG $(2, 15)$ or equivalent</p> <p>Use of $y=mx+c$ or $\frac{y-y_1}{x-x_1} = m$</p> <p>$y = -12x/10 + 17\frac{2}{5}$ or $y - 15 = -12/10(x - 2)$</p> <p>$6x + 5y - 87 = 0$ OR $-6x - 5y + 87 = 0$</p> | <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>A2</p> <p>12</p> | <p>Or equivalent. Allow 1 slip in sign of substitution Allow for sight of $10^2 + 12^2$</p> <p>CAO</p> <p>FT 'their FG' of equivalent difficulty expressed correctly, e.g. $\sqrt{104} = 2\sqrt{26}$, or $\sqrt{44} = 2\sqrt{11}$ needs to be in the form $a\sqrt{b}$ where $a \neq 1$ and $b \neq 1$ or simpler Sight of $2\sqrt{61}$ implies previous $\sqrt{244}$</p> <p>Or equivalent</p> <p>CAO. Mark final answer and then FT</p> <p>FT $-1/\text{grad FG}$</p> <p>Accept $(2, \dots)$ or $(\dots, 15)$</p> <p>CAO</p> <p>Must show substitution of 3 values Method to find the equation using mid-point and perpendicular gradient (not $10/12$ or $5/6$ or 'their gradient') FT their mid-point (not F or G) & their perpendicular gradient (not $10/12$ or $5/6$ or 'their gradient'), or FT substitution of their midpoint with their perpendicular gradient (not $10/12$ or $5/6$ or 'their gradient'), in $y = mx + c$ (towards finding c) <i>If no working for finding gradient is seen, then 'their 'spurious' incorrect perpendicular gradient' must be negative</i></p> <p>FT for correct unsimplified form, not written in quotient form, i.e. $\frac{y-15}{x-2} = \frac{-12}{10}$</p> <p>Accept terms in different orders provided '=0'</p> <p>CAO for A2 and A1</p> <p>A1 for $12x + 10y - 174 = 0$ or other multiple of the correct response (not with fractional coefficients), with integer values for a, b and c, and with terms in any order provided '=0', OR</p> <p>A1 for $5y = -6x + 87$ or equivalent correct simplified equation but not given in the required form</p> |

| | Additional Mathematics Summer 2017 | | Final Version |
|---|---|---|--|
| 7 | <p>(Arc length \Rightarrow) $2 \times \pi \times 5 \times 110/360$ $= 55\pi/18$ (cm)</p> <p>(Circumference) $55\pi/18 = 2 \times \pi \times$ cone radius or equivalent</p> <p>(Cone radius \Rightarrow) $55/36$ or $1.527(77\dots\text{cm})$</p> <p>(Perpendicular height² \Rightarrow) $5^2 -$ cone radius²</p> <p>(Perpendicular height \Rightarrow) $4.7608\dots(\text{cm})$</p> <p>(Volume of the cone \Rightarrow) $\frac{1}{3} \times \pi \times$ cone radius² \times perpendicular height</p> <p>(\Rightarrow) 11.6 (cm³)</p> <p>QWC2:</p> <ul style="list-style-type: none"> Candidates will be expected to present work clearly, with words explaining process or steps <p>AND</p> <ul style="list-style-type: none"> make few if any mistakes in mathematical form, spelling, punctuation and grammar in their answer <p>QWC1: Candidates will be expected to</p> <ul style="list-style-type: none"> present work clearly, with words explaining process or steps <p>OR</p> <ul style="list-style-type: none"> make few if any mistakes in mathematical form, spelling, punctuation and grammar in their final answer | <p>M1 A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>QWC 2</p> | <p>($= 9.599\dots\text{cm}$) May be implied in later working</p> <p>FT 'their derived arc length' Allow for $55\pi/18 = \pi \times$ diameter, if clearly diameter</p> <p>FT 'their derived cone radius', provided $\neq 5$</p> <p>FT 'their derived cone radius' and 'their derived perpendicular' provided neither value $\neq 5$</p> <p>May be shown in stages, e.g. method to find area first</p> <p>CAO. Award M1 only for $11.636\dots \text{cm}^3$ or 12cm^3</p> <p><i>Alternative (for 1st 4 marks) (not requiring arc length)</i> <i>(Area of sector \Rightarrow) $\pi \times 5^2 \times 110/360$ with intention to find the surface area of the cone</i> M1 $= 275\pi/36$ (cm²) ($=23.998\dots\text{cm}^2$) A1 <i>(Surface area of cone $= \pi rl$)</i> $275\pi/36 = \pi \times$ cone radius $\times 5$ M1 <i>(cone radius \Rightarrow) $55/36$ or $1.527(77\dots\text{cm})$</i> A1</p> <p>QWC2 Presents relevant material in a coherent and logical manner, using acceptable mathematical form, and with few if any errors in spelling, punctuation and grammar.</p> <p>QWC1 Presents relevant material in a coherent and logical manner but with some errors in use of mathematical form, spelling, punctuation or grammar OR evident weaknesses in organisation of material but using acceptable mathematical form, with few if any errors in spelling, punctuation and grammar.</p> <p>QWC0 Evident weaknesses in organisation of material, and errors in use of mathematical form, spelling, punctuation or grammar.</p> |
| 8 | <p>(a) $1140x^{18}$</p> <p>(b) $a = 3$ $b = 2$ $c = -6$</p> | <p>B2</p> <p>B1 B1 B1</p> <p>5</p> | <p>B1 for sight of $60x^{19}$. FT to 2nd B1 from $dy/dx = kx^n$ B0 for 60^{19} or 1140^{18} Ignore incorrect notation</p> <p><i>Accept sight of correct answers from 'uncorrected' working</i> <i>Do not accept embedded answers, candidates need to identify values for a, b and c, not accept as left in working without clearly stating.</i></p> |
| 9 | <p>$x = 5 \times \tan 60^\circ$ or $\tan 60^\circ = x/5$ ($x \Rightarrow$) $5\sqrt{3}$</p> | <p>M1 A1</p> <p>2</p> | <p>OR for use of $30^\circ, 60^\circ$ triangle with $5 \times$ sides of 1, 2 & $\sqrt{3}$ Do not award A1 unless a line of working is seen, i.e. $5\sqrt{3}$ without working is M0, A0</p> |

| | Additional Mathematics Summer 2017 | | Final Version |
|----|--|--|--|
| 10 | <p>(a) $10x^5/5 + 24x^3/3 - 2x + 3x^{-3}/-3$</p> <p style="text-align: right;">+ c (constant)</p> <p>(b) $12x^4/4 + 6x^3/3$</p> <p>$[12x^4/4 + 6x^3/3]^2_1$</p> <p style="text-align: center;">$= (3 \times 2^4 + 2 \times 2^3) - (3 \times 1^4 + 2 \times 1^3)$</p> <p style="text-align: center;">$= 59$</p> | <p>B4</p> <p>B1</p> <p>B2</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>10</p> | <p>B1 for each term ISW from correct unsimplified form. Simplified form is $2x^5 + 8x^3 - 2x - x^{-3}$</p> <p>Awarded only if at least B1 is awarded for integration</p> <p>Mark final answer, then FT. B1 for sight of $12x^4/4$ or $6x^3/3$ Ignore inclusion of '+c' throughout' (except final A mark)</p> <p>FT their integration, not original. Intention to use 2, 1 and subtract</p> <p>FT for correct use of limits Accept unsimplified fractions included</p> <p>CAO, not FT. Do not accept '59 + c' <i>Answer only, no working shown M0 A0 A0</i></p> |
| 11 | <p>$(dy/dx=) 9x^2 + 18x$ $dy/dx = 0$ or $9x^2 + 18x = 0$</p> <p style="text-align: center;">$x = 0$ and $y = 4$ $x = -2$ and $y = 16$</p> <p>$d^2y/dx^2 = 18x + 18$</p> <p>At (0, 4) $d^2y/dx^2 > 0$, point is a minimum At (-2, 16): $d^2y/dx^2 < 0$, point is a maximum</p> | <p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>7</p> | <p>FT their dy/dx form $ax^2 + bx$ throughout</p> <p><i>Answer only, no working shown M0 A0 A0</i></p> <p><i>Method for determining min or max MUST be shown, final answer only is M0 here, then A0,A0</i></p> <p>Or first derivative test, interpretation of first derivative test. Or alternative.</p> <p>FT for their x value (ignore y-values)</p> <p>FT for their other x value provided this does not have the same interpretation as the first x value (ignore y-values)</p> <p><i>SC1 for correct FT from $d^2y/dx^2 = ax + b, a > 0$, including allowed FT from $d^2y/dx^2 = 18x$, with $x = -2$ as maximum and $x = 0$ as a minimum (despite $d^2y/dx^2 = 0$)</i></p> <p><i>Do not accept trial & improvement methods unless both stationary points are found correctly and confirmed as stated in the mark scheme</i></p> |
| 12 | <p>$y + \delta y = (x + \delta x)^2 + 10(x + \delta x)$ Intention to subtract $(y =) x^2 + 10x$ to find δy $\delta y = 2x\delta x + (\delta x)^2 + 10\delta x$ Dividing by δx and (\lim) $\delta x \rightarrow 0$ $dy/dx = \lim_{\delta x \rightarrow 0} \delta y / \delta x = 2x + 10$</p> | <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>5</p> | <p>Or alternative notation. Allow if final bracket omitted</p> <p>Accept δx^2 as meaning $(\delta x)^2$</p> <p>FT equivalent level of difficulty CAO. Must follow from correct working <i>Use of dy/dx throughout or incorrect notation then possible maximum is only 4 marks, final A0</i></p> |
| 13 | <p>When $x = 4$, finding $y = 0$ $dy/dx = 10x - 20$</p> <p>(when $x = 4$) gradient is 20 Use of $\frac{y - y_1}{x - x_1} = m$ or $y = mx + c$</p> <p>$y - 0 = 20(x - 4)$ or $0 = 20 \times 4 + c, c = -80$ $20x - y - 80 = 0$</p> | <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>6</p> | <p>Ignore notation, e.g. $y = 10x - 20$, provided clear not from any wrong method.</p> <p>Method to form equation FT their y value but not $y = 4$, and their derived gradient</p> <p>CAO. Must be in this form, accept equivalents written as 3 terms with whole number coefficients with '=0' or '0='</p> |

| | Additional Mathematics Summer 2017 | | Final Version |
|----|--|---------------------------|---|
| 14 | $\frac{36}{2x+5} + \frac{55}{3x-1}$ <p>36(3x - 1) + 55(2x + 5) as a numerator</p> <p>(2x + 5)(3x - 1) as a denominator</p> $\frac{218x+239}{(2x+5)(3x-1)}$ | M2 A1 A1 A1 5 | <p>M1 for either fraction, or M1 for sight of $36 \div (2x + 5)$ or $55 \div (3x - 1)$, with or without brackets</p> <p>FT provided M1 awarded due to a slip in the second fraction</p> <p>FT provided M1 awarded due to a slip in the second fraction</p> <p>CAO. Mark final answer If the denominator is expanded it must be correct If no marks, award SC2 for an answer of $\frac{218x + 239}{1980}$ (from starting with inverted fractions)</p> |
| 15 | <p>(a) General sin curve intersecting x-axis only at (0°,0), (180°,0) and (360°,0)</p> <p>Correct curve with 4 and -4 on y-axis</p> <p>(b) 14.477... (°) and 165.522... (°) only</p> | M1 A1 B2 4 | <p>Allow general shape as the joining of key values, but straight rather than clearly curving towards a turn at 90° and 270° in particular</p> <p>Must show a clear curve, not straight at turning points</p> <p>Accept rounded or truncated, e.g. 14(°) with 166(°) or 15(°) with 165(°) These values need to be selected, not amongst others unless unambiguously indicated as the response. B1 for sight of 14.477... (°) or 165.522... (°) B0 for a pair of angles with sum 180°</p> |
| 16 | <p>(a) $(60)x^{16/8}/x^{1/5}$ or equivalent first stage of work evaluated correctly with simplification of indices</p> $60x^{9/5}$ <p>(b) Correctly extracting a factor of $x^{1/5}$ or $x^{3/5}$ (to give correct numerator) OR correct alternative method with one correct step towards simplification</p> $2x^{1/5} + x^{3/5}$ | B1 B1 M1 A1 4 | <p>CAO, must be simplified, allowing $60x^{14/5}$ Mark final answer</p> <p>For an alternative method award M1 for $2x^{1/5} + \dots$ or $\dots + x^{3/5}$</p> <p>CAO or equivalent factorised form. Mark final answer</p> |
| 17 | <p>(a) 225</p> <p>(b) $\frac{1}{8 + \sqrt{5}} \times \frac{8 - \sqrt{5}}{8 - \sqrt{5}}$</p> $= \frac{8 - \sqrt{5}}{59}$ | B1 M1 A1 3 | <p><i>No marks if no working. Must see $15^{1/5 \times 6}$ or 15^2 or $3^2 \times 5^2$ or 9×25</i></p> <p><i>No marks if no working.</i></p> <p>Mark final answer</p> |

Differentiating from first principles. Marking guide.

Q12.

| | | | |
|----|---|-------------------------------------|---|
| 12 | $y + \delta y = (x + \delta x)^2 + 10(x + \delta x)$ Intention to subtract $(y =) x^2 + 10x$ to find δy $\delta y = 2x\delta x + (\delta x)^2 + 10\delta x$ Dividing by δx and $(\lim) \delta x \rightarrow 0$ $dy/dx = \lim_{\delta x \rightarrow 0} \delta y/\delta x = 2x +$ 10 $\delta x \rightarrow 0$ | B1 M1 A1 M1 A1 5 | Or alternative notation. Allow if final bracket omitted Accept δx^2 as meaning $(\delta x)^2$ FT equivalent level of difficulty CAO. Must follow from correct working Use of dy/dx throughout or incorrect notation then possible maximum is only 4 marks, final A0 |
|----|---|-------------------------------------|---|

B1 For sight of $(x + \delta x)^2 + 10(x + \delta x)$ or $(x + h)^2 + 10(x + h)$ or using alternative notation. This mark is given whether $(x + \delta x)^2 + 10(x + \delta x)$ stands alone or is embedded in an expression or a formula.

M1 For the intent to subtract $x^2 + 10x$ from the above.

So $(x + \delta x)^2 + 10(x + \delta x) - x^2 + 10x$ will gain the M1 even though there are missing brackets.

It can also be awarded to those who have expanded $(x + \delta x)^2 + 10(x + \delta x)$ and then crossed out the x^2 term and the $+10x$ term.

Those who reverse the subtraction will gain M0 unless there is evidence later on of dividing by $-\delta x$.

A1 For sight of $2x\delta x + (\delta x)^2 + 10\delta x$ (Accept δx^2 as meaning $(\delta x)^2$) with no other terms. Treat as a CAO.

$2x + \delta x + 10$ will imply the above if division by δx has already been done.

M1 A FT, if of equivalent difficulty, is possible for this M1 (but not the subsequent A1).

A correct division by δx has to be done

(so if a FT it has to be correct for their $2x\delta x + (\delta x)^2 + 10\delta x$)

AND we must see 'lim $\delta x \rightarrow 0$ ' OR ' $\delta x \rightarrow 0$ ' OR ' δx tends to 0'.

It is M0 for ' $\delta x = 0$ ' OR ' $\delta x \approx 0$ ' OR ' δx is so small we can forget about it'.

All of the above marks can be gained even if there is no l.h.s. shown.

Final A1. Must be for a 'text book' quality presentation. E.g.

Has to be a correct l.h.s. for each line, ' δy ' or ' $\delta y/\delta x$ '

AND at some point ' $dy/dx = \lim_{\delta x \rightarrow 0} \delta y/\delta x$ ' or ' $dy/dx = \lim_{\delta x \rightarrow 0} 2x + \delta x + 10$ '