

Change to improper fractions first. Then get a common denominator.

Example: $2\frac{1}{5} - 1\frac{1}{2} = \frac{11}{5} - \frac{3}{2} = \frac{22}{10} - \frac{15}{10} = \frac{7}{10}$

1. $2\frac{1}{3} - 1\frac{1}{2}$

9. $2\frac{1}{5} - 1\frac{1}{3}$

2. $3\frac{1}{3} - 1\frac{3}{4}$

10. $4\frac{1}{2} - 1\frac{3}{4}$

3. $2\frac{1}{5} - 1\frac{1}{4}$

11. $3\frac{2}{5} - 1\frac{2}{3}$

4. $3\frac{2}{5} - 1\frac{3}{4}$

12. $4\frac{3}{5} - 2\frac{3}{4}$

5. $3\frac{1}{4} - 2\frac{2}{3}$

13. $5\frac{1}{5} - 2\frac{2}{3}$

6. $2\frac{1}{4} - 1\frac{4}{5}$

14. $3\frac{1}{10} - 2\frac{1}{3}$

7. $2\frac{2}{5} - 1\frac{2}{3}$

15. $1\frac{3}{10} - \frac{2}{3}$

8. $2\frac{1}{6} - 1\frac{1}{5}$

16. $2\frac{2}{5} - \frac{2}{3}$