

C2

Solutions and Mark Scheme

Final Version

1.	1	1.414213562		
	1.1	1.337908816		
	1.2	1.2489996		
	1.3	1.144552314	(5 values correct)	B2
	1.4	1.019803903	(3 or 4 values correct)	B1
		Correct formula with $h = 0.1$		M1
		$I \approx \frac{0.1}{2} \times \{1.414213562 + 1.019803903 + 2(1.337908816 + 1.2489996 + 1.144552314)\}$		
		$I \approx 0.494846946$		
		$I \approx 0.495$	(f.t. one slip)	A1
		Special case for candidates who put $h = 0.8$		
	1	1.414213562		
	1.08	1.35410487		
	1.16	1.286234815		
	1.24	1.209297317		
	1.32	1.121427662		
	1.4	1.019803903	(all values correct)	B1
		Correct formula with $h = 0.8$		M1
		$I \approx \frac{0.8}{2} \times \{1.414213562 + 1.019803903 + 2(1.35410487 + 1.286234815 + 1.209297317 + 1.121427662)\}$		
		$I \approx 0.495045871$		
		$I \approx 0.495$	(f.t. one slip)	A1

2. (a) $3 - 7 \cos \theta = 6(1 - \cos^2 \theta)$ (correct use of $\sin^2 \theta = 1 - \cos^2 \theta$) M1
 An attempt to collect terms, form and solve quadratic equation in $\cos \theta$, either by using the quadratic formula or by getting the expression into the form $(a \cos \theta + b)(c \cos \theta + d)$, with $a \times c =$ coefficient of $\cos^2 \theta$ and $b \times d =$ constant m1
 $6 \cos^2 \theta - 7 \cos \theta - 3 = 0 \Rightarrow (3 \cos \theta + 1)(2 \cos \theta - 3) = 0$
 $\Rightarrow \cos \theta = -\frac{1}{3},$ ($\cos \theta = \frac{3}{2}$) (c.a.o.) A1
 $\theta = 109.47^\circ, 250.53^\circ$ B1 B1
 Note: Subtract (from final two marks) 1 mark for each additional root in range from $3 \cos \theta + 1 = 0$, ignore roots outside range.
 $\cos \theta = -$, f.t. for 2 marks, $\cos \theta = +$, f.t. for 1 mark
- (b) $2x + 45^\circ = 35^\circ, 215^\circ, 395^\circ$ (one value) B1
 $x = 85^\circ, 175^\circ$ B1, B1
 Note: Subtract (from final two marks) 1 mark for each additional root in range, ignore roots outside range.
- (c) Correct use of $\tan \theta = \frac{\sin \theta}{\cos \theta}$ (o.e.) M1
 $\theta = 194.48^\circ$ A1
 $\theta = 345.52^\circ$ A1
 Note: Subtract (from final two marks) 1 mark for each additional root in range, ignore roots outside range.
3. (a) $x^2 = 8^2 + (x + 2)^2 - 2 \times 8 \times (x + 2) \times \cos 60^\circ$ (correct use of cos rule) M1
 $x^2 = 64 + x^2 + 4x + 4 - 8x - 16$ A1
 $x = 13$ (f.t. only $x = 21$ from $+ 16$ in the line above) A1
- (b) $\frac{\sin ACB}{8} = \frac{\sin 60^\circ}{13}$ (substituting correct values in the correct places in the sin rule, f.t. candidate's derived value for x) M1
 $ACB = 32.2^\circ$ (f.t. candidate's derived value for x) A1
4. (a) At least one correct use of the sum formula M1
 $\frac{8}{2} \times [2a + 7d] = 124$
 $\frac{20}{2} \times [2a + 19d] = 910$ (both correct) A1
 An attempt to solve the candidate's two equations simultaneously by eliminating one unknown M1
 $d = 5$ (c.a.o.) A1
 $a = -2$ (f.t. candidate's value for d) A1
- (b) $-2 + 5(n - 1) = 183$ (f.t. candidate's values for a and d) M1
 $n = 38$ (c.a.o.) A1

5. (a) $S_n = a + ar + \dots + ar^{n-1}$ (at least 3 terms, one at each end) B1
 $rS_n = ar + \dots + ar^{n-1} + ar^n$
 $S_n - rS_n = a - ar^n$ (multiply first line by r and subtract) M1
 $(1-r)S_n = a(1-r^n)$
 $S_n = \frac{a(1-r^n)}{1-r}$ (convincing) A1

(b) **Either:** $\frac{a(1-r^4)}{1-r} = 73 \cdot 8$
Or: $a + ar + ar^2 + ar^3 = 73 \cdot 8$ B1

$\frac{a}{1-r} = 125$ B1

An attempt to solve these equations simultaneously by eliminating one of the variables M1

$r^4 = 0.4096$ A1

$r = 0.8$ (c.a.o.) A1

$a = 25$ (f.t. candidate's value for r) A1

6. (a) $\frac{x^{4/3}}{4/3} - 2 \times \frac{x^{3/4}}{3/4} + c$ B1, B1

(-1 if no constant term present)

(b) (i) $5 + 4x - x^2 = 8$ M1

An attempt to rewrite and solve quadratic equation in x , either by using the quadratic formula or by getting the expression into the form $(x + a)(x + b)$, with $a \times b = 3$ m1
 $(x - 1)(x - 3) = 0 \Rightarrow x = 1, 3$ (c.a.o.) A1

Note: Answer only with no working earns 0 marks

(ii) **Either:**

$$\text{Total area} = \int_1^3 (5 + 4x - x^2) dx - \int_1^3 8 dx$$

(use of integration) M1

Either: $\int 5 dx = 5x$ **and** $\int 8 dx = 8x$ or: $\int 3 dx = 3x$ B1

$$\int 4x dx = 2x^2, \quad \int x^2 dx = \frac{x^3}{3}$$

B1 B1

$$\text{Total area} = [-3x + 2x^2 - (1/3)x^3]_1^3 \quad (\text{o.e.})$$

$$= (-9 + 18 - 9) - (-3 + 2 - 1/3)$$

(substitution of candidate's limits in at least one integral) m1

Subtraction of integrals with correct use of candidate's

x_A, x_B as limits m1

$$\text{Total area} = \frac{4}{3} \quad (\text{c.a.o.}) \text{ A1}$$

Or:

Area of rectangle = 16

(f.t. candidate's x -coordinates for A, B) B1

$$\text{Area under curve} = \int_1^3 (5 + 4x - x^2) dx$$

(use of integration) M1

$$= [5x + 2x^2 - (1/3)x^3]_1^3$$

(correct integration) B2

$$= (15 + 18 - 9) - (5 + 2 - 1/3)$$

(substitution of candidate's limits) m1

$$= \frac{52}{3}$$

Use of candidate's, x_A, x_B as limits and trying to find total area by subtracting area of rectangle from area under curve m1

$$\text{Total area} = \frac{52}{3} - 16 = \frac{4}{3} \quad (\text{c.a.o.}) \text{ A1}$$

7. (a) Let $p = \log_a x$
 Then $x = a^p$ (relationship between log and power) B1
 $x^n = a^{pn}$ (the laws of indices) B1
 $\therefore \log_a x^n = pn$ (relationship between log and power)
 $\therefore \log_a x^n = pn = n \log_a x$ (convincing) B1

(b) $\frac{1}{2} \log_a 324 = \log_a 324^{1/2}$
 $2 \log_a 12 = \log_a 12^2$ (at least one use of power law) B1

$\frac{1}{2} \log_a 324 + \log_a 56 - 2 \log_a 12 = \log_a \frac{324^{1/2} \times 56}{12^2}$

(use of addition law) B1

(use of subtraction law) B1

$\frac{1}{2} \log_a 324 + \log_a 56 - 2 \log_a 12 = \log_a 7$ (c.a.o) B1

Note: Answer only of $\log_a 7$ without any working earns 0 marks

(c) (i) $2^{x+1} = 2^x \times 2$ B1
 $3^x = 2^{x+1} \Rightarrow (1.5)^x = 2$ B1

(ii) **Hence:** $x \log_{10} 1.5 = \log_{10} 2$
 (taking logs on both sides and using the power law) M1
 (f.t. candidate's values for c and d)

$x = 1.71$ (c.a.o.) A1

Otherwise:

$x \log_{10} 3 = (x + 1) \log_{10} 2$
 (taking logs on both sides and using the power law) M1

$x = 1.71$ (c.a.o.) A1

8. (a) $A(-2, 4)$ B1
 A correct method for finding radius M1
 Radius = $\sqrt{10}$ A1

(b) An attempt to substitute $(3y - 4)$ for x in the equation of the circle M1
 $10y^2 - 20y + 10 = 0$ A1

Either: Use of $b^2 - 4ac$ m1
 Determinant = 0 $\Rightarrow x - 3y + 4 = 0$ is a tangent to the circle A1

Or: An attempt to factorise candidate's quadratic m1
 Repeated (single) root $\Rightarrow x - 3y + 4 = 0$ is a tangent to the circle A1

9. (a) $\frac{1}{2} \times 6^2 \times \sin \theta = 9.1$ M1
 $\theta = 0.53$ A1
- (b) Substitution of values in formula for area of sector M1
Area = $\frac{1}{2} \times 6^2 \times 0.53 = 9.54 \text{ cm}^2$ (f.t. candidate's value for θ) A1
- (c) $6 + 6 + 6\varphi = \pi \times 6$ M1
 $\varphi = 1.14$ A1
10. (a) $t_3 = 31$ B1
 $t_1 = 7$ (f.t. candidate's value for t_3) B1
- (b) All terms of the sequence are odd numbers E1