



GCE MARKING SCHEME

SUMMER 2018

**MATHEMATICS – M1 (LEGACY)
0980-01**

INTRODUCTION

This marking scheme was used by WJEC for the 2018 examination. It was finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conference was held shortly after the paper was taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conference, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about this marking scheme.

GCE MATHEMATICS – M1
SUMMER 2018 MARK SCHEME

Q	Solution	Mark	Notes
1(a)(i)	N2L on lift, upwards +ve $T - 1200g = 1200a$ $T = 1200(9.8 + 0.2)$ $T = 14160 \text{ (N)}$	M1 A1 A1	dim correct, all forces <i>T</i> and 1200 <i>g</i> opposing any correct form cao
1(a)(ii)	$T = 1200g \text{ (= 11760) (N)}$	B1	
1(b)	$Mg - R = Ma$ $M(9.8 - 3) = 442$ $M = 65$	M1 A1 A1	dim correct, all forces No extra any correct form cao


Q	Solution	Mark	Notes
2	Resolve in one direction $X = 16 - 9\cos 75^\circ - 21\sin 60^\circ$ $X = -4.5159$	M1 A1	obtain comp of resultant All forces, no extra
	Resolve in perpendicular direction $Y = 8 + 21\cos 60^\circ - 9\sin 75^\circ$ $Y = 9.8067$	M1 A1	obtain comp of resultant All forces, no extra.
	Resultant ² = $4.5159^2 + 9.8067^2$ Resultant = <u>10.8 (N)</u>	m1 A1	dep on both M's cao
	$\theta = \tan^{-1}\left(\frac{4.5159}{9.8067}\right)$ $\theta = \underline{24.7^\circ}$	m1 A1	 cao

Note

-1 if answers not 1 d.p.

Q	Solution	Mark	Notes
---	----------	------	-------

3.



Moments about P

$$36g \times 0.3 + R_Q \times 1.1 = 24g \times 0.7$$

$$R_Q = 53.45 \text{ (N)}$$

Resolve vertically

$$R_Q + R_P = 36g + 24g$$

$$R_P = 534.55 \text{ (N)}$$

M1	dim correct equation
B1	All forces, no extra
A1	any correct moment
A1	correct equation
A1	cao

M1	dim correct equation
A1	All forces, no extra
A1	cao

Notes

Moments about any point
 Correct moment
 Correct equation

M1	same conditions as above
B1	
A1	

Attempt at second equation
 Correct equation


M1	
A1	

Correct answers

A1A1	
------	--

Q Solution **Mark** **Notes**

4(a)



Apply N2L to Q

$$5g - T = 5a$$

Apply N2L to P

$$T - 3g = 3a$$

Adding

$$8a = 2g$$

$$a = \underline{2.45 \text{ (ms}^{-2}\text{)}}$$

$$T = \underline{36.75 \text{ N}}$$

M1 $5g$ and T opposing, dim. correct

A1 correct equ, allow $-ve a$

M1 $4g$ and T opposing, dim. Correct

A1 correct equ consistent with first equation

m1

A1 cao

A1 cao

4(b) Light string gives rise to tension constant throughout the string.

B1

4(c) Smooth peg means that the tensions in the strings on both sides of the peg are equal.

B1

Notes (Newton's method)

Attempt at Newton's method

M1 forces subtracted, masses added

$$5g - 3g = (5+3)a$$

A1

$$a = 2.45 \text{ (ms}^{-2}\text{)}$$

A1 cao

N2L applied to either particle

M1 wt and T opposing, dim. Correct

$$T - 3g = 3a$$

A1

$$T = 3(9.8 + 2.45)$$

m1 substitution

$$T = \underline{36.75 \text{ N}}$$


A1 cao

Q	Solution	Mark	Notes
5(a)	$I = \text{change in momentum}$ $I = 0.16(20 - (-12))$ $I = 5.12 \text{ (Ns)}$	M1 A1	used
5(b)	$I = Ft$ $5.12 = F \times \frac{1}{8}$ $F = 40.96 \text{ (N)}$	M1 A1	used ft answer in (a)

Q	Solution	Mark	Notes
6(a)	Vel of A when B starts to fall $v^2 = u^2 + 2as, u=0, a=(\pm)9.8, s=(\pm)0.1$ $v^2 = 0 + 2 \times 9.8 \times 0.1$ $v = \frac{7}{5}$	M1 A1 A1	oe complete method cao
6(b)	Vel of A when it reaches the ground $v^2 = u^2 + 2as, u=0, a=(\pm)9.8, s=(\pm)40$ $v^2 = 0 + 2 \times 9.8 \times 40$ $v = 28$	M1 A1	
6(c)	Time of travel of B = time for A to reach ground $v = u + at, u = \frac{7}{5}, v=28, a=9.8$ $28 = \frac{7}{5} + 9.8t$ $t = \frac{19}{7}$	M1 A1 A1	 ft (a) and (b)
	Distance travelled by B in that time $s = ut + \frac{1}{2}at^2, u=0, a=9.8, t=\frac{19}{7}$ $s = 0 + \frac{1}{2} \times 9.8 \times \left(\frac{19}{7}\right)^2$ $s = 36.1$	M1 A1 A1	 ft candidates' 19/7 cao
	Distance between A and $B = 40 - 36.1$ $= 3.9$ (m)	A1	cao

Q Solution**Mark Notes**

7(a)



Conservation of momentum

$$4m + 2m = mv + m \times 1.4v$$

$$2.4v = 6$$

$$v = 2.5$$

M1 dim correct equ.

A1

A1 cao

Restitution

$$1.4 \times 2.5 - 2.5 = -e(2 - 4)$$

$$e = 0.5$$

M1 no more than 1 sign error

A1 ft v in (a)A1 ft v in (a) provided $0 < e < 1$.7(b) Speed of B after collision = v'

$$v' = 3.5 \times 0.6$$

$$v' = 2.1 \text{ (ms}^{-1}\text{)}$$

M1 ft v_A A1 ft v_A 7(c) Distance between A and B at time of collision with the wall = $(3.5 - 2.5) \times 5$
= 5B1 ft $v, 1.4v$ After collision with wall, A and B

approach each other with

$$\text{velocity} = 2.1 + 2.5 = 4.6$$


B1 ft v, v' Time to second collision between A and B

$$= \frac{5}{4.6}$$

$$= 1.09 \text{ (s) (correct to 2 d.p.)}$$

B1 cao

Q	Solution	Mark	Notes
8.			



Resolve perpendicular to plane

$$R = 10g \cos \alpha$$

$$F = 10g\mu \cos \alpha$$

B1

With T acting upwards

N2L applied to particle

$$T - F - mg \sin \alpha = ma$$

$$98 - F - 10g \sin \alpha = 0$$

M1

dim correct, all forces

A1

With T acting downwards

N2L applied to particle

$$F - T' - 10g \sin \alpha = ma$$

$$F - 49 - 10g \sin \alpha = 0$$

M1

dim correct, all forces

A1

Adding

$$98 - 49 = 20 \times 9.8 \times \sin \alpha$$

$$\sin \alpha = \frac{1}{4}$$

$$\cos \alpha = \frac{\sqrt{15}}{4}$$

$$\mu = \frac{F}{R}$$

$$\mu = \frac{49 + 10 \times 9.8 \times 0.25}{10 \times 9.8 \times \frac{\sqrt{15}}{4}}$$

$$\mu = \frac{\sqrt{15}}{5} = \sqrt{\frac{3}{5}} = 0.7746$$

m1

A1

M1

A1

cao

Q	Solution	Mark	Notes
9(a).	$\bar{x} = 4$ (cm)	B1	
9(b)	Shape mass distance(y)		
	<i>ABCE</i> 40 2.5	B1	2.5
	<i>ECD</i> 36 8	B1	8
	<i>PQR</i> 12 7	B1	7
	<i>ABCDE</i> 64 \bar{y}	B1	areas
	Moments about <i>AB</i>	M1	dim correct equation
	$64\bar{y} + 12 \times 7 = 40 \times 2.5 + 36 \times 8$	A1	ft table if consistent
	$\bar{y} = 4.75$ (cm)	A1	cao