

C3

Solutions and Mark Scheme

Final Version

- 1.**
- | | | | |
|--|--|-------------|----------------------------|
| | 0 | 0.69314718 | |
| | 0.25 | 0.825939419 | |
| | 0.5 | 0.974076984 | |
| | 0.75 | 1.136871006 | (5 values correct) B2 |
| | 1 | 1.313261688 | (3 or 4 values correct) B1 |
| | Correct formula with $h = 0.25$ | | M1 |
| | $I \approx \frac{0.25}{3} \times \{0.69314718 + 1.313261688 + 4(0.825939419 + 1.136871006) + 2(0.974076984)\}$ | | |
| | $I \approx 11.80580453 \div 12$ | | |
| | $I \approx 0.983817044$ | | |
| | $I \approx 0.984$ | | (f.t. one slip) A1 |
-
- 2.** (a) e.g. $\theta = \frac{\pi}{2}$
- | | | | |
|--|---------------------------------------|---|----|
| | $\sin 4\theta = 0$ | (choice of θ and one correct evaluation) | B1 |
| | $4 \sin^3 \theta - 3 \sin \theta = 1$ | (both evaluations correct but different) | B1 |
- (b) $3(1 + \tan^2 \theta) = 7 - 11 \tan \theta$. (correct use of $\sec^2 \theta = 1 + \tan^2 \theta$) M1
- An attempt to collect terms, form and solve quadratic equation in $\tan \theta$, either by using the quadratic formula or by getting the expression into the form $(a \tan \theta + b)(c \tan \theta + d)$, with $a \times c =$ coefficient of $\tan^2 \theta$ and $b \times d =$ constant m1
- $3 \tan^2 \theta + 11 \tan \theta - 4 = 0 \Rightarrow (3 \tan \theta - 1)(\tan \theta + 4) = 0$
- $\Rightarrow \tan \theta = \frac{1}{3}, \tan \theta = -4$ (c.a.o.) A1
- $\theta = 18.4^\circ, 198.4^\circ$ B1
- $\theta = 104.0^\circ, 284.0^\circ$ B1 B1
- Note: Subtract 1 mark for each additional root in range for each branch, ignore roots outside range.
- $\tan \theta = +, -, \text{ f.t. for 3 marks, } \tan \theta = -, -, \text{ f.t. for 2 marks}$
- $\tan \theta = +, +, \text{ f.t. for 1 mark}$

3. (a) $\frac{d}{dx}(y^3) = 3y^2 \frac{dy}{dx}$ B1
 $\frac{d}{dx}(2x^3y) = 2x^3 \frac{dy}{dx} + 6x^2y$ B1
 $\frac{d}{dx}(3x^2 + 4x - 3) = 6x + 4$ B1
 $x = 2, y = 1 \Rightarrow \frac{dy}{dx} = -\frac{8}{19}$ (c.a.o.) B1

(b) (i) $\frac{dx}{dt} = 6t, \quad \frac{dy}{dt} = 12t^2 + 6t^5$ (all three terms correct) B2
(one term correct) B1

Use of $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1

$\frac{dy}{dx} = 2t + t^4$ (c.a.o.) A1

(ii) $\frac{d}{dt}\left[\frac{dy}{dx}\right] = 2 + 4t^3$ (f.t. candidate's expression for $\frac{dy}{dx}$) B1
 $\frac{d}{dx}$

Use of $\frac{d^2y}{dx^2} = \frac{d}{dt}\left[\frac{dy}{dx}\right] \div \frac{dx}{dt}$ M1

$\frac{d^2y}{dx^2} = \frac{1 + 2t^3}{3t}$ (c.a.o.) A1

4. $f(x) = 2 - 10x + \sin x$
An attempt to check values or signs of $f(x)$ at $x = 0, x = \pi/8$ M1
 $f(0) = 2 > 0, f(\pi/8) = -1.54 < 0$
Change of sign $\Rightarrow f(x) = 0$ has root in $(0, \pi/8)$ A1
 $x_0 = 0.2$
 $x_1 = 0.219866933$ (x_1 correct, at least 5 places after the point) B1
 $x_2 = 0.221809976$
 $x_3 = 0.221999561$
 $x_4 = 0.222018055 = 0.22202$ (x_4 correct to 5 decimal places) B1
An attempt to check values or signs of $f(x)$ at $x = 0.222015, x = 0.222025$ M1
 $f(0.222015) = 4.56 \times 10^{-5} > 0, f(0.222025) = -4.46 \times 10^{-5} < 0$ A1
Change of sign $\Rightarrow \alpha = 0.22202$ correct to five decimal places A1
Note: 'change of sign' must appear at least once

5. (a) $\frac{dy}{dx} = \frac{3}{1+(3x)^2}$ or $\frac{1}{1+(3x)^2}$ or $\frac{3}{1+3x^2}$ M1
 $\frac{dy}{dx} = \frac{3}{1+9x^2}$ A1
- (b) $\frac{dy}{dx} = \frac{ax+b}{2x^2-3x+4}$ (including $a=0, b=1$) M1
 $\frac{dy}{dx} = \frac{4x-3}{2x^2-3x+4}$ A1
- (c) $\frac{dy}{dx} = e^{2x} \times m \cos x + ke^{2x} \times \sin x$ ($m = \pm 1, k = 1, 2$) M1
 $\frac{dy}{dx} = e^{2x} \times m \cos x + ke^{2x} \times \sin x$ (either $m = 1$ or $k = 2$) A1
 $\frac{dy}{dx} = e^{2x} \times \cos x + 2e^{2x} \times \sin x$ (c.a.o.) A1
- (d) $\frac{dy}{dx} = \frac{(1+\cos x) \times m \sin x - (1-\cos x) \times k \sin x}{(1+\cos x)^2}$ ($m = \pm 1, k = \pm 1$) M1
 $\frac{dy}{dx} = \frac{(1+\cos x) \times -(-\sin x) - (1-\cos x) \times (-\sin x)}{(1+\cos x)^2}$ A1
 $\frac{dy}{dx} = \frac{2 \sin x}{(1+\cos x)^2}$ A1

6. (a) (i) $\int \frac{1}{4x-7} dx = k \times \ln|4x-7| + c \quad (k = 1, 4, 1/4)$ M1
 $\int \frac{1}{4x-7} dx = 1/4 \times \ln|4x-7| + c$ A1

(ii) $\int e^{3x-1} dx = k \times e^{3x-1} + c \quad (k = 1, 3, 1/3)$ M1
 $\int e^{3x-1} dx = 1/3 \times e^{3x-1} + c$ A1

(iii) $\int \frac{5}{(2x+3)^4} dx = -\frac{5}{3k} \times (2x+3)^{-3} + c \quad (k = 1, 2, 1/2)$ M1
 $\int \frac{5}{(2x+3)^4} dx = -\frac{5}{6} \times (2x+3)^{-3} + c$ A1

(b) $\int \sin\left[2x + \frac{\pi}{4}\right] dx = \left[k \times \cos\left[2x + \frac{\pi}{4}\right] \right] \quad (k = -1, -2, \pm 1/2)$ M1

$\int \sin\left[2x + \frac{\pi}{4}\right] dx = \left[-1/2 \times \cos\left[2x + \frac{\pi}{4}\right] \right]$ A1

$\int_0^{\pi/4} \sin\left[2x + \frac{\pi}{4}\right] dx = k \times \left[\cos\left[\frac{3\pi}{4}\right] - \cos\left[\frac{\pi}{4}\right] \right]$
(f.t. candidate's value for k) A1

$\int_0^{\pi/4} \sin\left[2x + \frac{\pi}{4}\right] dx = \frac{\sqrt{2}}{2}$ (c.a.o.) A1


7. (a) $2|x+1| - 3 = 7 \Rightarrow |x+1| = 5$ B1
 $x = 4, -6$ B1

(b) Trying to solve either $5x - 8 \geq 3$ or $5x - 8 \leq -3$ M1
 $5x - 8 \geq 3 \Rightarrow x \geq 2.2$
 $5x - 8 \leq -3 \Rightarrow x \leq 1$ (both inequalities) A1
Required range: $x \leq 1$ or $x \geq 2.2$ (f.t. one slip) A1

Alternative mark scheme

$(5x - 8)^2 \geq 9$ (forming and trying to solve quadratic) M1
Critical points $x = 1$ and $x = 2.2$ A1
Required range: $x \leq 1$ or $x \geq 2.2$ (f.t. one slip in critical points) A1


8.



- The x -axis is an asymptote for $f(x)$ at $-\infty$, correct behaviour at $+\infty$ M1
- $y = f(x)$ cuts y -axis at $(0, 1)$ A1
- $y = 2f(x) - 3$ cuts y -axis at $(0, -1)$ (f.t. candidate's y -intercept for $f(x)$) B1
- $y = -3$ is an asymptote for $2f(x) - 3$ at $-\infty$, with graph above $y = -3$ B1
- The diagram shows that the graph of $y = 2f(x) - 3$ is steeper than the graph of $y = f(x)$ in the first quadrant B1

9. (a) $y = \frac{1}{2}\sqrt{x} - 3$ and an attempt to isolate x M1
 $2y = \sqrt{x} - 3 \Rightarrow x = 4y^2 + 3$ A1
 $f^{-1}(x) = 4x^2 + 3$ (f.t. one slip in candidate's expression for x) A1
 $R(f^{-1}) = [4, \infty)$ B1
 $D(f^{-1}) = [1/2, \infty)$ B1

(b)



- $y = f^{-1}(x)$ a parabola B1
starting at $(1/2, 4)$ (f.t. candidate's $D(f^{-1})$) B1
 $y = f(x)$ as in diagram (c.a.o.) B1

10. (a) $R(f) = (-1, \infty)$ B1
 $R(g) = (3, \infty)$ B1
- (b) $f(1) = 0$ is not in the domain of g E1
- (c) (i) $fg(x) = (2x - 1)^2 - 1$ M1
 $fg(x) = 4x(x - 1)$ or $4x^2 - 4x$ A1
(ii) $D(fg) = (2, \infty)$ B1
 $R(fg) = (8, \infty)$ B1