

C3

1. (a)
- | | | | | |
|--|---|-------------|-------------------------|----|
| | 1 | 1.386294361 | | |
| | 1.25 | 1.517870719 | | |
| | 1.5 | 1.658228077 | | |
| | 1.75 | 1.802122256 | (5 values correct) | B2 |
| | 2 | 1.945910149 | (3 or 4 values correct) | B1 |
| | Correct formula with $h = 0.25$ | | | M1 |
| | $I \approx \frac{0.25}{3} \times \{1.386294361 + 1.945910149 + 4(1.517870719 + 1.802122256) + 2(1.658228077)\}$ | | | |
| | $I \approx 19.92863256 \div 12$ | | | |
| | $I \approx 1.66071938$ | | | |
| | $I \approx 1.6607 \quad \text{(f.t. one slip)}$ | | | |

Note: Answer only with no working earns 0 marks

(b)

$$\int_1^2 \ln \left[\frac{1}{3+x^2} \right] dx \approx -1.6607 \quad \text{(f.t. candidate's answer to (a))} \quad \text{B1}$$

2. $2 \operatorname{cosec}^2 \theta + 3(\operatorname{cosec}^2 \theta - 1) + 4 \operatorname{cosec} \theta = 9$
 (correct use of $\cot^2 \theta = \operatorname{cosec}^2 \theta - 1$) M1

An attempt to collect terms, form and solve quadratic equation in $\operatorname{cosec} \theta$, either by using the quadratic formula or by getting the expression into the form $(a \operatorname{cosec} \theta + b)(c \operatorname{cosec} \theta + d)$, with $a \times c =$ coefficient of $\operatorname{cosec}^2 \theta$ and $b \times d =$ candidate's constant m1

$$5 \operatorname{cosec}^2 \theta + 4 \operatorname{cosec} \theta - 12 = 0 \Rightarrow (5 \operatorname{cosec} \theta - 6)(\operatorname{cosec} \theta + 2) = 0$$

$$\Rightarrow \operatorname{cosec} \theta = \frac{6}{5}, \operatorname{cosec} \theta = -2$$

$$\Rightarrow \sin \theta = \frac{5}{6}, \sin \theta = -\frac{1}{2} \quad \text{(c.a.o.)} \quad \text{A1}$$

$$\theta = 56.44^\circ, 123.56^\circ \quad \text{B1}$$

$$\theta = 210^\circ, 330^\circ \quad \text{B1 B1}$$

Note: Subtract 1 mark for each additional root in range for each branch, ignore roots outside range.

$\sin \theta = +, -$, f.t. for 3 marks, $\sin \theta = -, -$, f.t. for 2 marks

$\sin \theta = +, +$, f.t. for 1 mark

3. (a) $\frac{d(2x^3)}{dx} = 6x^2$, $\frac{d(2x)}{dx} = 2$, $\frac{d(25)}{dx} = 0$ B1
 $\frac{d(x^2 \cos y)}{dx} = x^2(-\sin y) \frac{dy}{dx} + 2x(\cos y)$ B1
 $\frac{d(y^4)}{dx} = 4y^3 \frac{dy}{dx}$ B1
 $\frac{dy}{dx} = \frac{6x^2 + 2x \cos y + 2}{x^2 \sin y - 4y^3}$ (c.a.o.) B1
- (b) (i) candidate's x -derivative = $3t^2$
candidate's y -derivative = $4t + 20t^3$
(one term correct B1, all three terms correct B2)
 $\frac{dy}{dx} = \frac{\text{candidate's } y\text{-derivative}}{\text{candidate's } x\text{-derivative}}$ M1
 $\frac{dy}{dx} = \frac{4 + 20t^2}{3t}$ (c.a.o.) A1
- (ii) $\frac{dy}{dx} = 5 \Rightarrow 20t^2 - 15t + 4 = 0$
(f.t. candidate's expression for $\frac{dy}{dx}$ from (i)) B1
Considering $b^2 - 4ac$ for candidate's quadratic M1
 $b^2 - 4ac = 225 - 320 < 0$ and hence no such real value of t exists
(f.t. candidate's quadratic) A1
4. (a) $f'(x) = (11) \times g(x) - 6x$
where $g(x) = \text{either } \frac{2}{1 + (2x)^2} \text{ or } \frac{1}{1 + (2x)^2} \text{ or } \frac{2}{1 + 2x^2}$ M1
 $f'(x) = 11 \times \frac{2}{1 + 4x^2} - 6x$ A1
 $f'(x) = 0 \Rightarrow 12x^3 + 3x - 11 = 0$ (convincing) A1
- (b) $x_0 = 0.9$
 $x_1 = 0.884366498$ (x_1 correct, at least 5 places after the point) B1
 $x_2 = 0.886029122$
 $x_3 = 0.885852598$
 $x_4 = 0.885871344 = 0.88587$ (x_4 correct to 5 decimal places) B1
Let $h(x) = 12x^3 + 3x - 11$
An attempt to check values or signs of $h(x)$ at $x = 0.885865$,
 $x = 0.885875$ M1
 $h(0.885865) = -1.42 \times 10^{-4} < 0$, $h(0.885875) = 1.70 \times 10^{-4} > 0$ A1
Change of sign $\Rightarrow \alpha = 0.88587$ correct to five decimal places A1

5. (a) $\frac{dy}{dx} = \frac{1}{3} \times (9 - 2x)^{-2/3} \times f(x)$ $(f(x) \neq 1)$ M1
 $\frac{dy}{dx} = \frac{-2}{3} \times (9 - 2x)^{-2/3}$ A1
- (b) $\frac{dy}{dx} = \frac{f(x)}{\cos x}$ (including $f(x) = 1$) M1
 $\frac{dy}{dx} = \frac{\pm \sin x}{\cos x}$ A1
 $\frac{dy}{dx} = -\tan x$ (c.a.o.) A1
- (c) $\frac{dy}{dx} = x^3 \times f(x) + \tan 4x \times g(x)$ M1
 $\frac{dy}{dx} = x^3 \times f(x) + \tan 4x \times g(x)$
(either $f(x) = 4 \sec^2 4x$ or $g(x) = 3x^2$) A1
 $\frac{dy}{dx} = x^3 \times 4 \sec^2 4x + \tan 4x \times 3x^2$ (all correct) A1
- (d) $\frac{dy}{dx} = \frac{(3x + 2)^4 \times k \times e^{6x} - e^{6x} \times 4 \times (3x + 2)^3 \times m}{[(3x + 2)^4]^2}$
with either $k = 6, m = 3$ or $k = 6, m = 1$ or $k = 1, m = 3$ M1
 $\frac{dy}{dx} = \frac{(3x + 2)^4 \times 6 \times e^{6x} - e^{6x} \times 4 \times (3x + 2)^3 \times 3}{[(3x + 2)^4]^2}$ A1
 $\frac{dy}{dx} = \frac{18x \times e^{6x}}{(3x + 2)^5}$ (c.a.o.) A1

6. (a) (i) $\int \frac{9}{4x+3} dx = k \times 9 \times \ln|4x+3| + c$ ($k = 1, 4, 1/4$) M1
 $\int \frac{9}{4x+3} dx = 9/4 \times \ln|4x+3| + c$ A1
- (ii) $\int 3e^{5-2x} dx = k \times 3 \times e^{5-2x} + c$ ($k = 1, -2, -1/2$) M1
 $\int 3e^{5-2x} dx = -3/2 \times e^{5-2x} + c$ A1
- (iii) $\int \frac{5}{(7x-1)^3} dx = \frac{k \times 5 \times (7x-1)^{-2}}{-2} + c$ ($k = 1, 7, 1/7$) M1
 $\int \frac{5}{(7x-1)^3} dx = \frac{5 \times (7x-1)^{-2}}{-2 \times 7} + c$ A1


Note: The omission of the constant of integration is only penalised once.

- (b) $\int \cos\left[3x - \frac{\pi}{6}\right] dx = \left[\frac{k \times \sin\left[3x - \frac{\pi}{6}\right]}{\left[\frac{\pi}{6} \right]} \right]$ ($k = 1, 3, \pm 1/3$) M1
- $\int \cos\left[3x - \frac{\pi}{6}\right] dx = \left[\frac{1/3 \times \sin\left[3x - \frac{\pi}{6}\right]}{\left[\frac{\pi}{6} \right]} \right]$ A1
- $\int_0^{\pi/3} \cos\left[3x - \frac{\pi}{6}\right] dx = k \times \left[\frac{\sin\left[\frac{5\pi}{6}\right] - \sin\left[-\frac{\pi}{6}\right]}{\left[\frac{\pi}{6} \right]} \right]$
- (A correct method for substitution of limits
f.t. only candidate's value for k , $k = 1, 3, \pm 1/3$) m1
- $\int_0^{\pi/3} \cos\left[3x - \frac{\pi}{6}\right] dx = \frac{1}{3}$ (c.a.o.) A1

Note: Answer only with no working earns 0 marks

7. (a) Choice of a, b , with one positive and one negative and one side correctly evaluated M1
Both sides of identity evaluated correctly A1
- (b) Trying to solve $2x + 1 = 3x - 4$ M1
Trying to solve $2x + 1 = -(3x - 4)$ M1
 $x = 5, x = 0.6$ (both values) A1
- Alternative mark scheme**
- $(2x + 1)^2 = (3x - 4)^2$ (squaring both sides) M1
 $5x^2 - 28x + 15 = 0$ (c.a.o.) A1
 $x = 5, x = 0.6$ (both values, f.t. one slip in quadratic) A1

8.



- Correct shape, including the fact that the y -axis is an asymptote for $y = f(x)$ at $-\infty$ B1
 $y = f(x)$ cuts x -axis at $(1, 0)$ B1
 Correct shape, including the fact that $x = -3$ is an asymptote for $y = \frac{1}{2}f(x+3)$ at $-\infty$ B1
 $y = \frac{1}{2}f(x+3)$ cuts x -axis at $(-2, 0)$ (f.t. candidate's x -intercept for $f(x)$) B1
 The diagram shows that the graph of $y = f(x)$ is steeper than the graph of $y = \frac{1}{2}f(x+3)$ in the first quadrant B1

9. (a) $y + 3 = e^{2x+1}$ B1
 An attempt to express equation as a logarithmic equation and to isolate x M1
 $x = \frac{1}{2} [\ln(y+3) - 1]$ (c.a.o.) A1
 $f^{-1}(x) = \frac{1}{2} [\ln(x+3) - 1]$ (f.t. one slip in candidate's expression for x) A1
- (b) $D(f^{-1}) = (a, b)$ with B1
 $a = -3$ B1
 $b = -2$ B1

10. (a) $R(f) = (-19, \infty)$ B1
 $R(g) = (-\infty, -2)$ B1
- (b) $D(fg) = (6, \infty)$ B1
 $R(fg) = (-15, \infty)$ B1
- (c) (i) $fg(x) = \left[1 - \frac{1}{2}x \right]^2 - 19$ B1
- (ii) Putting expression for $fg(x)$ equal to $2x - 26$ and setting up a quadratic in x of the form $ax^2 + bx + c = 0$ M1
 $\frac{1}{4}x^2 - 3x + 8 = 0 \Rightarrow x = 4, 8$ (c.a.o.) A1
4
Rejecting $x = 4$ and thus $x = 8$ (c.a.o.) A1